


STAR as a Fixed Target Experiment?



Brooke Haag

Hartnell College / University of California, Davis


Presented at the Meeting of the California Division of the APS

November 13, 2009






The Basics



matter in the universe is made of atoms



nucleus = protons + neutrons


THE STANDARD MODEL

	Fermions			Bosons	
Quarks	<i>u</i> up	<i>c</i> charm	<i>t</i> top	γ photon	Force carriers
	<i>d</i> down	<i>s</i> strange	<i>b</i> bottom	<i>Z</i> Z boson	
Leptons	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	<i>W</i> W boson	
	<i>e</i> electron	μ muon	τ tau	<i>g</i> gluon	
	Higgs boson*				

*Yet to be confirmed

Source: AAAS



mesons = 2 quarks
baryons = 3 quarks




nucleons are hadrons (made of quarks)

Creating mini-big bangs in the laboratory


Goal: Use relativistic collisions of nuclei to create hot dense matter which reproduces the earliest stages of the universe




1. Ions about to collide*



2. Ion collision




3. Quarks, gluons freed



4. Plasma created

QCD phase diagram



- We have discovered the QGP !
- We have not located the critical point.
- A beam energy scan will explore regions of lower temperature and higher chemical potential to find the critical point.
- Can we extend the physics analysis at lower energies by introducing a target into the STAR detector ?



Kinematic Calculations

Collision Energy (GeV)	Single Beam Energy	Single Beam P _z (GeV/c)	Fixed Target √s	Single Beam Rapidity	Center of Mass Rapidity
22.4 Cu+Cu	11.2	11.16	4.66 Cu+Al	3.18	1.59
19.6 Au+Au	9.8	9.76	4.47 Au+Al	3.04	1.52
9.2 Au+Au	4.6	4.50	3.21 Au+Al	2.28	1.14

√s_{NN} = center of mass energy


- $\sqrt{s_{NN}} = \sqrt{2m^2 + 2Em}$
m = 0.9315 GeV/c² ; E = 9.8 GeV
- $\sqrt{s_{NN}} = 4.47$ GeV

- $p_z = \sqrt{E^2 - m^2} = 9.76$ GeV/c

rapidity (y)

- $y_{beam} = 0.5 * [\ln(E + p_z) / (E - p_z)]$
- $y_{beam} = 3.0$
- $y_{cm} = 1.5$

STAR detector array






Event Selection





Cu+Al Event Cuts:

- $|zVertex| > 75$ cm
- $rVertex > 2$ cm
- multiplicity > 11
- $\Sigma p_z * zVertex < 0$




Spectra



- Fits to spectra assume Boltzmann distributions
- With efficiency corrections we can extract a temperature for comparison with previous measurements by other experiments

Next Steps

- We need to understand the acceptance of the detector in the forward region
- One important issue -- looking in the forward region with the TPC for particle identification





Conclusions and Outlook

- We can do physics with STAR as a fixed target experiment !
 - We have been able to extract spectra from several species for fixed target collisions at lab rapidity
 - Will it contribute to the critical point search?
 - need to understand detector efficiency at high rapidities
 - need much better statistics - this study is a proof of principle
 - want to get yields and slopes which compare favorably with published data in this energy range